

IIT-HEP-95/5

hep-ex/9510002

PRODUCTION OF CHARM, CHARMONIUM, AND BEAUTY IN 800 GEV PROTON-NUCLEON COLLISIONS ^a

DANIEL M. KAPLAN ^b
*Illinois Institute of Technology,
Chicago, IL 60616, USA*

for the Fermilab E789 Collaboration ^c

Fermilab Experiment 789 took data during the 1991 fixed-target run on the production of charm, charmonium, and beauty in 800 GeV proton-nucleon collisions. Several results have been published or submitted for publication; only a brief synopsis will be given here. The apparatus is shown schematically in Figs. 1 and 2; it is based on the pre-existing E605/772 spectrometer, augmented for this run primarily by the addition of two silicon-microstrip vertex telescopes (Fig. 2), one above and one below the beam. In most events only two oppositely-charged particle tracks traversing the entire spectrometer are reconstructed, one passing through each telescope.




Figure 1: E789 apparatus (plan view).

E789 results include the nuclear dependences of charmonium production, which goes as $A^{\alpha(x_F)}$ with $\alpha(0.05) = 0.90 \pm 0.02$,^{1,2} and of $D^0(\overline{D}^0)$ production,

^ato appear in *Proc. 6th Int. Conf. on Hadron Spectroscopy (Hadron 95)*, Manchester, England, 10–14 July 1995.

^bE-Mail address: kaplan@fnal.gov.

^cC. N. Brown, T. A. Carey, Y. C. Chen, R. Childers, W. E. Cooper, C. W. Darden, G. Gidal, K. N. Gounder, P. M. Ho, L. D. Isenhower, D. M. Jansen, R. G. Jeppesen, D. M. Kaplan, J. S. Kapustinsky, G. C. Kiang, M. S. Kowitt, D. W. Lane, L. M. Lederman, M. J. Leitch, J. W. Lillberg, W. R. Luebke, K. B. Luk, P. L. McGaughey, C. S. Mishra, J. M. Moss, J. C. Peng, R. S. Preston, D. Pripstein, J. Sa, M. E. Sadler, R. Schnathorst, M. H. Schub, V. Tanikella, P. K. Teng, J. R. Wilson; Abilene Christian University, Academia Sinica (Taiwan), University of California at Berkeley, University of Chicago, Fermi National Accelerator Laboratory, Illinois Institute of Technology, Lawrence Berkeley Laboratory, National Cheng Kung University, Northern Illinois University, University of South Carolina.




Figure 2: E789 vertex telescopes (elevation view).

which depends linearly on target atomic weight ($\alpha = 1.02 \pm 0.03 \pm 0.02$ at $x_F \simeq 0.03$).³ The lower values of α observed for charmonium (especially for x_F near zero) are consistent with models in which charmonium production is suppressed in nuclei due to dissociation by comovers.² Models in which the nuclear suppression of charmonium production is an initial-state effect (*e.g.* due to possible nuclear modification of the gluon structure function) are disfavored, since they would predict similar nuclear dependences for J/ψ and D production.

We have measured differential cross sections for charm³ and charmonium⁴ production. The observed D^0 cross section is $d\sigma/dx_F = 58 \pm 3 \pm 7 \mu\text{b}/\text{nucleon}$, which extrapolated over all x_F implies a total D^0 cross section $\sigma = 17.7 \pm 0.9 \pm 3.4 \mu\text{b}/\text{nucleon}$.³ Averaging with previous measurements using 800 GeV proton beams⁵ gives $\sigma(pN \rightarrow D^0 X) + \sigma(pN \rightarrow \bar{D}^0 X) = (20.9 \pm 3.5) \mu\text{b}/\text{nucleon}$, consistent with next-to-leading-order (NLO) QCD predictions⁶ within the broad range of theoretical uncertainty. However, as at the Tevatron Collider,⁷ charmonium production is substantially underestimated, at least in models which include only contributions from color-singlet charmonium states below $D\bar{D}$ threshold.¹⁰ We find $\sigma(p + N \rightarrow J/\psi + X) = 442 \pm 2 \pm 88 \text{ nb}/\text{nucleon}$ and $\sigma(p + N \rightarrow \psi' + X) = 75 \pm 5 \pm 22 \text{ nb}/\text{nucleon}$, factors of 7 and 25 above QCD predictions.⁴

Production of beauty hadrons is studied by searching for evidence of $J/\psi \rightarrow \mu^+ \mu^-$ decay occurring in vacuum downstream of the 3-mm-long Au target. A significant excess is observed of events with vertex downstream of the target compared to those with vertex upstream, leading to the measured cross section for J/ψ from b decay $d^2\sigma/dx_F dp_T^2 = 107 \pm 28 \pm 19 \text{ pb}/(\text{GeV}/c)^2/\text{nucleon}$ at $x_F = 0.05$ and $p_T = 1 \text{ GeV}/c$.⁸ This can be corrected for the $b \rightarrow J/\psi + X$ branching ratio and extrapolated over all of phase

space to yield $\sigma(pN \rightarrow b\bar{b} + X) = 5.7 \pm 1.5 \pm 1.3 \text{ nb/nucleon}$.⁸ This value is consistent with NLO QCD predictions but a factor ≈ 2 below their central value.⁶

The small beauty cross section at fixed-target energy is surprising given that the beauty cross section at the Tevatron collider is a factor 2 *above* the central QCD prediction.⁹ It appears difficult to accomodate this steep s dependence within the next-to-leading-order model. However, caution is in order since the assumption of factorization of b -quark production and fragmentation may be unreliable in hadroproduction, especially at the low p_T values of the fixed-target data.⁶

Considering that the range of QCD predictions included 3 nb,⁶ our measurement should be taken as favorable for the HERA- B project, as is also our success in reconstructing $b \rightarrow J/\psi + X$ decay vertices in a fixed-target environment at a 50 MHz interaction rate. This latter accomplishment also indicates the feasibility of a fixed-target charm experiment with sensitivity ≈ 2000 times higher than that of existing experiments, for which a Letter of Intent is in progress.¹¹

References

1. M. S. Kowitt *et al.*, Phys. Rev. Lett. **72**, 1318 (1994).
2. M. J. Leitch *et al.*, "Nuclear Dependence of J/ψ Production by 800 GeV/ C Protons near $x_F = 0$," to appear in Phys. Rev. D (1995).
3. M. J. Leitch *et al.*, Phys. Rev. Lett. **72**, 2542 (1994).
4. M. H. Schub *et al.*, Phys. Rev. D **52**, 1307 (1995).
5. R. Ammar *et al.*, Phys. Rev. Lett. **61**, 2185 (1988); K. Kodama *et al.*, Phys. Lett. **263B**, 573 (1991).
6. M. L. Mangano *et al.*, Nucl. Phys. **B405**, 507 (1993); S. Frixione *et al.*, Nucl. Phys. **B431**, 453 (1994).
7. E. Braaten and S. Fleming, Phys. Rev. Lett. **74**, 3327 (1995).
8. D. M. Jansen *et al.*, Phys. Rev. Lett. **74**, 3118 (1995).
9. F. Abe *et al.*, Fermilab-Conf-94/136-E; F. Abe *et al.*, Phys. Rev. D **50**, 4252 (1994); S. Abachi *et al.*, Phys. Rev. Lett. **74**, 3548 (1995).
10. For discussion of alternative charmonium-production models see *e.g.* P. R. Page, Nucl. Phys. **B446**, 189 (1995) and references therein.
11. D. M. Kaplan, this Conference ("Future Facilities" session); D. M. Kaplan, "High-Impact Charm Physics at the Turn of the Millennium," IIT-HEP-95/3, hep-ex/9508018, to appear in *Proc. Tau-Charm Factory Workshop*, Argonne National Laboratory, June 21–23, 1995, and references therein.